

# NSW Immunisation Schedule

from 1 July 2018



AGE	DISEASE	VACCINE
<b>CHILDHOOD VACCINES</b>		
Birth	Hepatitis B	H-B-VAX II <b>OR</b> ENGERIX B
6 weeks	Diphtheria, tetanus, pertussis, <i>Haemophilus influenzae</i> type b, hepatitis B, polio	INFANRIX HEXA
	Pneumococcal	PREVENAR 13
	Rotavirus	ROTARIX
4 months	Diphtheria, tetanus, pertussis, <i>Haemophilus influenzae</i> type b, hepatitis B, polio	INFANRIX HEXA
	Pneumococcal	PREVENAR 13
	Rotavirus	ROTARIX
6 months <sup>†</sup>	Diphtheria, tetanus, pertussis, <i>Haemophilus influenzae</i> type b, hepatitis B, polio	INFANRIX HEXA
12 months	Meningococcal ACWY	NIMENRIX
	Pneumococcal	PREVENAR 13
	Measles, mumps, rubella	MMR II <b>OR</b> PRIORIX
18 months	Diphtheria, tetanus, pertussis	INFANRIX <b>OR</b> TRIPACEL
	Measles, mumps, rubella, varicella	PRIORIX TETRA <b>OR</b> PROQUAD
	<i>Haemophilus influenzae</i> type b	ACT-HIB
4 years	Diphtheria, tetanus, pertussis, polio	INFANRIX-IPV <b>OR</b> QUADRACEL
<b>ADOLESCENT VACCINES - SCHOOL VACCINATION PROGRAM</b>		
Year 7	Diphtheria, tetanus, pertussis	BOOSTRIX
	Human papillomavirus (2 doses)	GARDASIL 9
Years 10 - 11 (In 2018)	Meningococcal ACWY	MENACTRA
<b>ADULT VACCINES</b>		
Pregnant women	Influenza (Annually-any trimester)	INFLUENZA
	Pertussis (Third trimester, ideally 28-32 weeks)	BOOSTRIX <b>OR</b> ADACEL
65 years and over	Influenza (Annually)	FLUAD <b>OR</b> FLUZONE HIGH DOSE
	Pneumococcal (One dose)*	PNEUMOVAX 23
70 years (Catch-up for 71-79 years until 31 October 2021)	Zoster	ZOSTAVAX
<b>AT RISK GROUPS</b>		
6 months and over with medical risk conditions <sup>†</sup>		INFLUENZA
All children 6 months to < 5 years (In 2018)	Influenza (annual)	
Aboriginal people 15 years and over		
Aboriginal people 15-49 years with medical risk factors		
Aboriginal people 50 years and over	Pneumococcal*	PNEUMOVAX 23
65 years and over		

<sup>†</sup> Refer to the current online edition of The Australian Immunisation Handbook for all medical risk factors and conditions

\* Refer to the current edition of The Australian Immunisation Handbook for timing of doses

<sup>‡</sup> at risk children require an additional dose of pneumococcal (Prevenar 13)